

Grewinium, a substitute name for *Grewioxylon* Shallom non Schuster

RASHMI SRIVASTAVA AND J.S. GULERIA

Birbal Sahni Institute of Palaeobotany, 53 University Road, Lucknow 226 007, India.
Email: rashmi_bsip@yahoo.com

(Received 24 January 2000; revised version accepted 05 June 2000)

SCHUSTER (1910) instituted a fossil wood genus *Grewioxylon* (Type: *G. swedenborgii*) showing affinities with modern *Grewia laevigata* of family Tiliaceae. Kräuse (1922) transferred *Grewioxylon swedenborgii* to *Dipterocarpoxylon* Holden (1916) belonging to family Dipterocarpaceae on account of the presence of vertical gum canals. Moreover, absence of *Pterospermum* type of tile cells in *G. swedenborgii* justifies this transfer. Subsequently Schweitzer (1958) transferred this species under *Shoreoxylon* Den Berger (1923) as *S. swedenborgii* due to presence of vertical gum canals in concentric rings in it. Shallom (1963), ignorant of Schuster's (1910) publication, again instituted the genus *Grewioxylon* for the fossil wood of *Grewia*. Although this fact was recognized by Prakash and Dayal (1965) who, however, maintained Schuster's genus because of its priority and emended its diagnosis. It would have been better that Prakash and Dayal and subsequent workers had placed their species under the genus *Grewioxylon* Shallom, instead of *Grewioxylon* Schuster, as Schuster's genus had already became an illegitimate genus vide Art. 56.1 of International Code of Botanical Nomenclature (Greuter *et al.*, 1994) due to transfer of its type species to other genus (Greuter *et al.*, 1994).

Since *Grewioxylon* Shallom, became a later homonym of *Grewioxylon* Schuster, as it is spelt exactly like the name based on different type that was previously validly published for the taxon of same rank as per Art. 53.1 of ICBN. Therefore, it is considered desirable to propose a substitute name (*nomen novum*) according to Art. 58.3 of ICBN. Hence, a substitute name *Grewinium* is proposed for *Grewioxylon* Shallom (1963) non Schuster (1910).

GREWINIUM Srivastava & Guleria nom. nov.

GREWIOXYLON Shallom, 1963, p. 171

Type species—*Grewinium intertrappeum* Srivastava & Guleria comb. nov., J. Indian bot. Soc. 42 : 171, pl. 1, figs 1-5, text figs A-F; 1963.

Extended Diagnosis—Wood-diffuse-porous. Growth rings-indistinct or delimited by narrow line of terminal parenchyma. Vessels-small to medium sized; solitary and in radial multiples of 2-4 (or more); vessel members short with oblique or tailed ends; perforations simple; intervessel pits small, bordered and alternate. Parenchyma-paratracheal present, terminal and diffuse present or absent; strands irregularly storied. Rays-fine to broad; heterocellular; narrow rays have storied tendency; broad rays consist of *Pterospermum* type of tile cells interspersed among procumbent cells. Radial gum canal like structures absent or present in median portion of broad rays. Fibres-nonseptate, irregularly storied.

Remarks—Extended diagnosis is given because of addition of certain anatomical characters like radial gum canal type structures are present/absent in some Indian Deccan Intertrappean (Maastrichtian-Danian) woods, which differentiate *Grewinium* from extant *Grewia*. Although radial gum canals are not of common occurrence in the family Tiliaceae, Mouton and Jacquet (1981) have recorded their sporadic occurrence in some genera.

Hence fossil woods of *Grewioxylon* except two informal species, viz., *Grewioxylon* sp. cf. *G. mahurzariense* Lakhanpal *et al.*, 1978 and *Grewioxylon* sp. Selmeier, 1985 described until now are named as:

Grewinium intertrappeum Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon intertrappea* Shallom, J. Indian bot. Soc. 42 : pl. 1, figs 1-5, text figs A-F; 1963.

Grewinium indicum Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon indicum* Prakash & Dayal, Palaeobotanist 13: 21, pl. 1, figs 6, 7, pl. 2, figs 8-13, text fig. 2; 1965.

Grewinium mahurzariense Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon malurzariense* Prakash & Dayal, Palaeobotanist 13 : 17, pl. 1, figs 1-5, text fig. 1; 1965.

Grewinum fontanesii Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon fontanesii*, Vozentin-Serra, Palaeontographica 177B: 146, pl. 5, figs 1-5, pl. 6, figs 1-4; 1981.

Grewinum neumaieri Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon neumaieri* Selmeier, Münchner Geowiss. Abh. 6: 98, pl. 3, figs 1-3, pl. 4, fig. 2, pl. 5, figs 3, 5, pl. 8, figs 1, 2; 1985.

Grewinum ortenburgense Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon ortenburgense* Selmeier, Münchner Geowiss. Abh. 6: 102, pl. 3, fig. 4, pl. 4, figs 3, 4, pl. 5, figs 1, 2, pl. 6, figs 1-4, pl. 7, figs 1, 2, 4, pl. 8, figs 3-6; 1985.

Grewinum canalisum Srivastava & Guleria, comb. nov.

Basionym: *Grewioxylon canalisum* Bande & Srivastava, Geophytology 24: 131, pl. 1, figs 1-7; 1995.

Acknowledgements—The authors are grateful to Prof Anshu K Sinha, Director, Birbal Sahni Institute of Palaeobotany, Lucknow, for granting permission to publish the paper (BSIP/RCPC/PUBL/2000-5). They are also grateful to the referees for their valuable suggestions in improving the manuscript.

REFERENCES

- Bande MB & Srivastava R 1995. *Grewia*-type of woods from the Deccan Intertrappean beds of India. Geophytology 24 : 131-135.
 Den Berger LG 1923. Fossiele houtsoorten uit het Tertiair van Zuid-Sumatra. Geologie en mijnbouw 7 : 143-148.

Greuter W, Barrie FR, Burdet HM, Chaloner WG, Demoulin V, Hawksworth DL, Jørgensen PM, Nicolson DH, Silva PC, Trehane P & McNeil J 1994. International Code of Botanical Nomenclature (Tokyo Code), adopted by the Fifteenth International Botanical Congress, Yokohama, August-September 1993. Regnum Vegetabile 131.

Holden R 1916. A fossil wood from Burma. Records of the Geological Survey of India. 47 : 267-272.

Kräusel R 1922. Fossile Hölzer aus dem Tertiär von Süd Sumatra. Verhandelingen van het Geologisch-Mijnbouwkundig Genootschap voor Nederland en Koloniën, Geologische Series 5 : 231-287.

Lakhanpal RN, Prakash U & Bande MB 1978. Fossil dicotyledonous woods from the Deccan Intertrappean beds of Mandla District in Madhya Pradesh. Palaeobotanist 25 : 190-204.

Mouton J & Jacquet P 1981. Contribution à L'inventaire floristique et xylologique de la Guyane Française - 7 - *Apetiba*, *Luehea*, *Lueheopsis* (Tiliaceae). 106th Congress national des Societe's savantes, Perpignan, Science, Fasc. II : 245-259.

Prakash U & Dayal R 1965. Fossil woods of *Grewia* from the Deccan Intertrappean Series, India. Palaeobotanist 13 : 17-24.

Schuster J 1910. Über Nicolien und Nicolien ähnliche Hölzer. Kungliga Svenska vetenskapsakademiens handlingar 45 : 1-18.

Schweitzer HJ 1958. Die fossilen Dipterocarpaceen- Hölzer Palaeontographica 105B : 1-166.

Selmeier A 1985. Jungtertiäre Kieselhölzer (Fam. Bombacaceae, Tiliaceae, Meliaceae) aus dem Ortenburger Schotter von Rauscheröd, Niederbayern. Münchner Geowissenschaftliche 6 : 89-140.

Shallom LJ 1963. A fossil dicotyledonous wood with tile cells, from the Deccan Intertrappean beds of Mahurzari. Journal of the Indian Botanical Society 42 : 170-176.

Vozentin-Serra C 1981. Les structures ligneuses néogènes du plateau de Di Linch (Sud-Viêtnam). Palaeontographica 177B : 136-161.