

Palynofloral, palaeovegetational and palaeoenvironmental investigations in the Lower Kamthi Formation of South India

HARINAM JOSHI¹, NEHA AGGARWAL^{*1}, NEERJA JHA¹

Birbal Sahni Institute of Palaeosciences, 53 University Road, Lucknow – 226 007, India.

*Email: neha_264840@yahoo.co.in**

ABSTRACT

These studies were carried out to better understand the palynostratigraphic and palaeoclimatic fluctuations observed in the floral ecosystem. The palynofloral investigation discovered two palynoassemblages (I-II). *Faunipollenites* spp. and *Striatopodocarpites* spp. dominate Palynoassemblage-I (430-232.10 m), with a high incidence of *Striasulcites* spp. Palynoassemblage-II (208.30-83.50 m) is distinguished by striate bisaccates and a high *Densipollenites* spp. frequency. *Alisporites* sp., *Falcisporites nuthaliensis*, *Klausipollenites schaubergeri*, *Chordasporites australiensis*, *Guttulapollenites hannonicus*, and *Corisaccites alutus* are the younger elements of these palynoassemblages. Guadalupian (Wordian-Capitanian) and Lopingian (Wuchiapingian-Changhsingian) ages have been assigned to the palynoassemblage-I and II based on palynofloral evidence. Organic matter in various forms indicates four distinct palynofacies assemblages (PF I-IV). According to their findings, the sequence is dominated by the presence of sub-arborescent/arborescent forest cover that thrived in swampy settings near the depositional site. During deposition, the host sediments exhibit oxic to anoxic conditions as well as variable energy levels of the freshwater regime.

Keywords- Palynology, Palynofacies, Guadalupian, Lopingian, Godavari Valley Coalfield.

INTRODUCTION

The Godavari Valley Coalfield is a NNW-SSE trending rift basin, located between N 16°38' to N 19°32' and E 79°12' to E 81°39'. The gravity anomalies show a large rift valley where Gondwana sediments were deposited in a block faulted trough that was gradually formed (Qureshy *et al.*, 1968). Later glaciations (post-Barakar Formation deposits) are suggested to have caused the Godavari Coalfield to first emerge as a Sag-type basin before changing it into a rift basin (Biswas, 1999). Based on the tectonics and structure Godavari Coalfield has been divided into four sub-basins *viz.* (i) Godavari Sub-basin, (ii) Kothagudem Sub-basin, (iii) Chintalapudi Sub-basin, and (iv) Krishna–Godavari Coastal Tract. In the current work, the subsurface sedimentary sequence in borehole MJR-13 from the Jangareddygudem region of the Chintalapudi Sub-basin is discussed.

Several boreholes in the Chintalapudi Sub-basin have been analysed by various authors for palynological studies (Bottapagudem: Jha, 2004; Amavaram: Srivastava & Jha, 1992; Sattupalli: Jha, 2008; Srivastava & Jha, 1994; Ayyanpalli-Gompana: Srivastava & Jha, 1993; Gattugudem: Jha, 2002). Jha *et al.* have recently published a review on the palynostratigraphy of the Chintalapudi Sub-basin (2018). Considered to be younger and predominantly developed during the Kamthi period is the Chintalapudi sub-basin, largely due to the general absence of the Barakar and Barren Measures formations over a significant part of it. Except for Upper Barakar and Barren Measures, a complete palynofloral succession from the Early Permian to Early Triassic sequence has been documented. However, all previous studies focused solely on the palynostratigraphic setup in the basin, whereas the current study includes palynofacies investigation as well as palynostratigraphy, which is useful for establishing the palaeoenvironmental and depositional settings. Very few studies are available from the Chintalapudi sub-basin in context to the palaeoenvironmental setup (Mishra *et al.*, 2018; Mishra and Singh, 2018; Aggarwal *et al.*, 2019).

In this communication, borehole MJR-13 from the Jangareddygudem area, located on the sub-northeastern basin's margin, has been rendered for palynological studies aimed at understanding the stratigraphy, depositional environment, and palynofloral transition. Comparing coeval sequences from intrabasinal and interbasinal correlations in India and other Gondwana continents with contemporaneous deposits has also yielded age estimates. Singareni Collieries Company Limited (SCCL) of South India drilled borehole MJR-13 in the Jangareddygudem area. Figure 1 shows the location map for borehole MJR-13 in detail. Palynofacies studies are used to better understand the palaeodepositional settings in the study area. Palynofacies analysis reveals information about the depositional environment in the study area as well as the maturation of organic matter. Tyson (1993, 1995) and Batten (1996) classified dispersed organic matter, and the current study follows their classification for palaeoenvironmental studies. The objective of this study is to utilize palynoflora for correlating and dating sediments, alongside investigating palaeodepositional environments within the Chintalapudi sub-basin. A total of 37 samples, sourced from different lithologies within the Kamthi Formation, were collected and subjected to analysis for this purpose.

GEOLOGICAL SETTINGS OF THE AREA

Godavari Coalfield deposits from the Permian, Triassic, and Jurassic periods, as well as the Lower Cretaceous, span the entire Gondwana (Asselian-Albian) period in the South Indian Peninsula (Mukhopadhyay *et al.*, 2010). Lower Gondwana (Asselian- Changhsingian) sediments are exposed along the basin's eastern and western margins in this linear belt. The Upper Gondwana (Induan-Albian) sediments, on the other hand, cover the central/axial portion of the basin.

The Chintalapudi Sub-basin, which extends the Kothagudem Sub-basin to the southeast (Fig. 1), is unconformably situated atop the Archean gneisses and spans an area of 2500 square kilometres. The formations known as the Talchir, Barakar, Kamthi, Maleri, Kota, and

Gangapur represent the stratigraphic succession of the Gondwana sediments in this Sub-basin (Table 1). The overlaying sequence in this sub-basin encompasses the axial portion of the basin, while the Talchir and Barakar formations are exposed at the eastern and western basin edges. In the past, the Gondwana rocks of this sub-basin have been described as the Chintalapudi Formation, Kamthi Formation, and Kamthi Sandstone (Blanford, 1872; Rao, 1982; Raiverman *et al.*, 1985). The Chintalapudi Sub-basin was formerly said to be of a younger generation and to have grown mostly during the deposition of the Kamthi, as evidenced by the widespread lack of the Barakar and Barren Measures formations over a significant portion of the sub-basin (Raiverman *et al.*, 1985). Later, the stratigraphy was corrected by Lakshminarayana and Murti (1990) and Lakshminarayana (1996), who also noticed the existence of the Upper Gondwana sequence and discovered the Talchir and Barakar formations. As a result, the Kamthi Formation likewise sits on top of the Barakar Formation on the southwest edge of the Chintalapudi Sub-basin. Still, it is faulted against the foundation rock further south. Barring Barren Measures, the palynological study of the 19 boreholes in this sub-basin (Jha *et al.*, 2018) shows that the Early Permian–Early Triassic succession is nearly complete.




Figure 1. Study area map of Godavari Graben with location map of the borehole MJR-13 in the Jangareddygudem area.

The stratigraphy of the Kamthi Formation in the Chintalapudi Sub-basin remains poorly understood and warrants further investigation (refer to Table 1). This necessitates the execution of dating and correlation efforts in this area, which is the focus of the present study.

Table 1. A generalized stratigraphical sequence in Chintalapudi Sub-basin (after Lakshminarayana, 1996).

Age	Group	Formation	Lithology	
EARLY CRETACEOUS	UPPER GONDWANA	Gangapur	White sandstone, buff siltstone and claystone.	
JURASSIC		Unconformity		
		Kota	UPPER-sandstone and conglomeratic sandstone MIDDLE- limestone and marl LOWER-Conglomeratic sandstone and trough cross stratified sandstone.	
		Unconformity		
		Maleri	(?) Red clay and lime-pellet rock	
MIDDLE TRIASSIC-LATE TRIASSIC		UPPER GONDWANA	Upper Kamthi	Coarse-grained, ferruginous sandstones with clay clasts and pebbles and subordinate violet cherty siltstones and pebble beds.
EARLY TRIASSIC			Middle Kamthi	Alternating sequence of medium-grained white to greenish grey white sandstones and buff to greenish grey clays.
			Unconformity	
LATE PERMIAN	LOWER G	Lower Kamthi/ Raniganj	Medium-to-coarse-grained, greyish-white sandstones with a few coal seams	
		Unconformity		

EARLY PERMIAN	O N D W A N A	Barakar	UPPER: White feldspathic sandstone, siltstone, shale, carbonaceous shale and coal seams. LOWER: Very coarse-grained, pebbly, feldspathic sandstone
		Talchir	Diamictite, rhythmite, fine-grained light green sandstone, and siltstone
Unconformity			
PRECAMBRIAN	-----	Proterozoic	
		Archaean	

MATERIAL AND METHODS


Palynofloral analysis: The thirty-seven rock samples from various lithologies were collected from the drilled borehole MJR-13 (Fig. 2 & 3) in the Kamthi Formation. The running borehole was drilled up to 477.50 m, and no workable coal seam has been marked as per lithological information. The thin coal layers identified in the sequence belong to the Kamthi Formation. Samples were processed by standard palynological maceration techniques proposed by Bharadwaj (1962) and Bharadwaj and Salujha (1964).

Palynofacies analysis: A standard non-oxidative procedure (Batten, 1999; Prasad *et al.*, 2013; Aggarwal *et al.*, 2019) proceeded to recover palynofacies matter. The residue was only mounted on slides after treatment with hydrochloric acid (HCL) and hydrofluoric acid (HF).

The chemically treated materials for both analysis were sieved, and slides were prepared using Canada balsam as a mounting medium. For stratigraphic analysis, two hundred (200) palynomorphs and 500 organic matter particles were counted per sample for palynofacies analysis. The optical examination was conducted utilizing an Olympus BX61

microscope paired with a DP-25 camera and Cell A software. All slides are housed at the Museum, Birbal Sahni Institute of Palaeosciences, Lucknow.

ACCEPTED



15 **Figure 2.** Lithological column and sample levels, stratigraphic occurrence, and relative abundance of various palynomorphs in borehole MJR-

16 13.

PALYNOFLORAL INVESTIGATION

In the complete sequence, two distinct palynological assemblages have been identified, encompassing 49 genera of spores and pollen. Among these, one spore genus belongs to Sphenophyta and five to Pteridophyta. The Gymnosperms are represented by six monosaccate genera, ten striate bisaccate genera, fourteen non-striate bisaccate genera, five taeniate genera, one monosulcate, one praecolpate genus, and two costate genera along with three algal genera and one with unknown botanical affinity. The list of palynomorph taxa with their probable botanical affinity is given in Table 2. The recovered pollen grains range from well-preserved to broken states. Stratigraphically significant taxa have been illustrated in Figures 4, 5, and 6. The stratigraphic range and relative abundance of palynomorphs in borehole MJR-13 are illustrated in Figures 2.

Table 2. *List of spores and pollen grains with their probable botanical affinities* (Balme, 1995; Costamagna et al., 2018).

Botanical Affinity (Balme, 1995; Costamagna et al., 2018)	Recovered palynoflora from the studied borehole MJR-13
	PTERIDOPHYTES
	(i) Sphenophyta
Equisetopsida	<i>Calamospora</i> (= <i>Punctatisporites</i>) <i>exile</i> Bharadwaj & Salujha 1964
Equisetopsida	<i>Calamospora</i> (= <i>Punctatisporites</i>) sp.
	(ii) Filicophyta
Filicopsid	<i>Horriditriletes ramosus</i> (Balme & Hennelly) Bharadwaj & Salujha 1964
Filicopsid	<i>Horriditriletes</i> sp.
Filicopsid	<i>Lacinitriletes</i> sp.
Filicopsid	<i>Leiotriletes</i> sp.
Filicopsid	<i>Lophotriletes rectus</i> Bharadwaj & Salujha 1964
Filicopsid	<i>Lophotriletes</i> sp.
Filicopsid	<i>Microfoveolatispora foveolata</i> Tiwari & Singh 1986
	GYMNOSPERMS
	(i) Monosulcate grains
Cycadales	<i>Cycadopites</i> sp.
	(ii) Praecolpate grains
Unknown	<i>Praecolpatites nidpurensis</i> Bharadwaj & Srivastava 1969
	(iii) Monosaccate grains
Cordaitales	<i>Caheniasaccites elongatus</i> Bose & Kar 1966
Cordaitales	<i>Caheniasaccites</i> sp.
Unknown	<i>Kamthisaccites kamthiensis</i> Srivastava & Jha 1986
Cordaitales	<i>Parasaccites</i> (= <i>Cannanaropollis</i>) <i>densicarpus</i> Lele 1975
Cordaitales	<i>Parasaccites diffusus</i> (Tiwari) Bose & Maheshwari 1968
Cordaitales	<i>Parasaccites distinctus</i> Tiwari 1965
Cordaitales	<i>Parasaccites korbaensis</i> Bharadwaj & Tiwari 1964
Cordaitales	<i>Parasaccites obscures</i> Tiwari 1965
Cordaitales	<i>Parasaccites</i> sp.
Cordaitales	<i>Plicatipollenites indicus</i> Lele 1963
Cordaitales	<i>Plicatipollenites ovatus</i> Kar 1968
Cordaitales	<i>Plicatipollenites</i> sp.
Conifers	<i>Potoniéisporites</i> sp.
Coniferales	<i>Striomonosaccites ovatus</i> Bharadwaj 1962
	(iv) Bisaccate grains
	(a) Nonstriate bisaccates
Conifers	<i>Alisporites indarraensis</i> Segroves 1969
Conifers	<i>Alisporites landianus</i> Balme 1970

Conifers	<i>Alisporites</i> sp.
Unknown	<i>Brachysaccus</i> sp.
Peltaspermales	<i>Chordasporites australiensis</i> de Jersey 1962
Unknown	<i>Cuneatisporites</i> sp.
Conifers	<i>Falcisporites nuthaliensis</i> (Clark) Balme 1970
Conifers	<i>Falcisporites</i> sp.
Conifers	<i>Falcisporites zapfei</i> (Potonié & Klaus) Leschik 1956
Glossopteridales	<i>Ibisporites diplosaccus</i> Tiwari 1968
Peltaspermales	<i>Klausipollenites schaubergeri</i> (Potonié & Kremp) Klaus 1963
Peltaspermales	<i>Klausipollenites</i> sp.
Peltaspermales	<i>Platysaccus</i> sp.
Conifers	<i>Primuspollenites levis</i> Tiwari 1964
Conifers	<i>Primuspollenites</i> sp.
Unknown	<i>Rhizomaspora indica</i> Tiwari 1965
Conifers	<i>Scheuringipollenites barakarensis</i> (Tiwari) Tiwari 1973
Conifers	<i>Scheuringipollenites maximus</i> (Hart) Tiwari 1973
Conifers	<i>Scheuringipollenites</i> sp.
Conifers	<i>Scheuringipollenites tentulus</i> (Tiwari) Tiwari 1973
Unknown	<i>Staurosaccites</i> sp.
Unknown	<i>Vesicaspora luteus</i> Salujha 1965
Unknown	<i>Vestigisporites rudis</i> Balme & Hennelly 1955
	(b) Striate bisaccates
Glossopterids	<i>Crescentipollenites barakarensis</i> Sinha 1972
Glossopterids	<i>Crescentipollenites densus</i> Bharadwaj & Srivastava 1969
Glossopterids	<i>Crescentipollenites globosus</i> (Maithy) Jha 1996
Glossopterids	<i>Crescentipollenites</i> sp.
Glossopteridales	<i>Faunipollenites bharadwajii</i> Maheshwari 1967
Glossopteridales	<i>Faunipollenites enigmatus</i> Maheshwari 1969
Glossopteridales	<i>Faunipollenites goraiensis</i> (Potonié & Lele) Maithy (1965)
Glossopteridales	<i>Faunipollenites parvus</i> Tiwari 1965
Glossopteridales	<i>Faunipollenites singrauliensis</i> Sinha 1972
Glossopteridales	<i>Faunipollenites</i> (=Protohaploxypinus) sp.
Glossopteridales	<i>Faunipollenites varius</i> (Bharadwaj) Tiwari <i>et al.</i> 1989
Coniferales	<i>Hamiapollenites insolitus</i> Bharadwaj & Salujha 1964
Unknown	<i>Lahirites</i> sp.
Glossopteridales	<i>Striapollenites saccates</i> Bharadwaj 1962
Coniferales	<i>Striasulcites ovatus</i> Venkatachala & Kar 1968
Coniferales	<i>Striasulcites</i> sp.
Coniferales	<i>Striasulcites tectus</i> Venkatachala & Kar, 1968
Coniferales	<i>Striatites communis</i> Bharadwaj & Salujha 1964
Coniferales	<i>Striatites parvus</i> Tiwari 1965
Coniferales	<i>Striatites reticuloidus</i> Tiwari 1965
Coniferales	<i>Striatites rhombicus</i> Bharadwaj & Salujha 1964

Coniferales	<i>Striatites</i> sp.
Coniferales	<i>Striatites tentulus</i> Tiwari 1965
Glossopteridales	<i>Striatopodocarpites brevis</i> Sinha 1972
Glossopteridales	<i>Striatopodocarpites cancellatus</i> (Balme & Hennelly 1955) Hart 1963
Glossopteridales	<i>Striatopodocarpites diffusus</i> Bharadwaj & Salujha 1964
Glossopteridales	<i>Striatopodocarpites magnificus</i> Bharadwaj & Salujha 1964
Glossopteridales	<i>Striatopodocarpites multistriatus</i> Jha 1996
Glossopteridales	<i>Striatopodocarpites</i> sp.
Glossopteridales	<i>Striatopodocarpites subcircularis</i> Sinha 1972
Glossopteridales	<i>Striatopodocarpites tiwarii</i> Bharadwaj & Dwivedi 1981
Glossopterids	<i>Strotersporites communis</i> (Wilson 1962) Hart 1964
Glossopterids	<i>Strotersporites</i> sp.
Glossopterids	<i>Strotersporites wilsonii</i> Klaus 1963
Glossopterids	<i>Strotersporites crassiletus</i> Jha 1996
Glossopterids	<i>Strotersporites perfectus</i> Maheshwari 1966
Glossopterids	<i>Verticipollenites debilis</i> Venkatachala & Kar 1968
Glossopterids	<i>Verticipollenites secretus</i> Bharadwaj 1962
Glossopterids	<i>Verticipollenites finimitus</i> Bharadwaj & Salujha 1964
Glossopterids	<i>Verticipollenites</i> sp.
	(c) Taeniate bisaccates
Coniferales	<i>Corisaccites alutus</i> Venkatachala & Kar 1968
Unknown	<i>Crucisaccites</i> sp.
Coniferales	<i>Guttulapollenites gondwanensis</i> Goubin 1965
Coniferales	<i>Guttulapollenites hannonicus</i> Goubin 1965
Coniferales	<i>Guttulapollenites</i> sp.
Unknown	<i>Lueckisporites virkkiae</i> Potonié & Klaus 1954
Coniferales	<i>Lunatisporites ovatus</i> (Goubin) Maheshwari & Banerji 1975
Coniferales	<i>Lunatisporites pellucidus</i> (Goubin) Helby 1972
Coniferales	<i>Lunatisporites</i> sp.
	(d) Costate grains
Coniferales	<i>Tiwariasporis</i> sp.
Coniferales	<i>Weylandites circularis</i> Bharadwaj & Srivastava 1969
Coniferales	<i>Weylandites indicus</i> Bharadwaj & Srivastava 1969
Coniferales	<i>Weylandites Lucifer</i> (Bharadwaj & Salujha) Bharadwaj & Dwivedi 1981
	ALGAE
	<i>Inaperturopollenites</i> sp.
	<i>Maculatasporites</i> sp.
	<i>Latosporites colliensis</i> (Balme & Hennelly) Bharadwaj 1962
	<i>Latosporites</i> sp.
	Unknown Botanical Affinity
	<i>Densipollenites densus</i> Bharadwaj & Srivastava 1969

	<i>Densipollenites indicus</i> Bharadwaj 1962
	<i>Densipollenites invisus</i> Bharadwaj & Salujha 1964
	<i>Densipollenites magnicarpus</i> Tiwari & Rana 1981
	<i>Densipollenites magnicarpus</i> var. <i>annulatus</i> Jha 1996
	<i>D. magnicarpus</i> var. <i>distinctus</i> Jha 1996
	<i>Densipollenites marginalis</i> Jha 1996
	<i>Densipollenites</i> sp.

ACCEPTED




Figure 3. Stratigraphically significant Palynotaxa of palynoassemblage-I (Wordian- Capitanian) of borehole MJR-13

A. *Faunipollenites singrauliensis*, BSIP Slide No. 15328, P57/3, B. *Striatopodocarpites magnificus*, BSIP Slide No. 15329, K48/3, C. *Faunipollenites enigmatus*, BSIP Slide No. 15325, K49, D. *F. singrauliensis*, MJR-13, BSIP Slide No. 15328, U57/4, E. *Strotersporites* sp., BSIP Slide No. 15329, P45/4, F. *Strotersporites crassiletus*, BSIP Slide No. 15326, M44/3, G. *Striatites reticuloidus*, BSIP Slide No. 15328, P32/4, H. *Striatopodocarpites* sp., BSIP Slide No. 15329, K55/2, I. *Striasulcites tectus*, BSIP Slide No. 15329, G62/1, J. *S. tectus*, BSIP Slide No. 15332, E62, K. *S. tectus*, BSIP Slide No. 15329, U54/4, L. *S. tectus*, BSIP Slide No. 15328, H38/4, M. *Faunipollenites singrauliensis*, BSIP Slide No. 15326, S33/3, N. *Faunipollenites* sp., BSIP Slide No. 15325, K35/2, O. *Crescentipollenites densus*, BSIP Slide No. 15327, H46, P. *Verticipollenites finimitus*, BSIP Slide No. 15322, H51/4,




Figure 4. Stratigraphically significant Palynotaxa of palynoassemblage-II (Wuchiapingian-Changhsingian) of borehole MJR-13

A. *Densipollenites magnicarpus*, BSIP Slide No. 15321, W57/4, B. *D. invisus*, BSIP Slide No. 15325, J52, C. *D. invisus*, BSIP Slide No. 15325, J52/4, D. *D. invisus*, BSIP Slide No. 15325, K37/3, E. *D. indicus*, BSIP Slide No. 15320, E61/1, F. *D. marginalis*, BSIP Slide No. 15332, P53/4, G. *D. magnicarpus*, BSIP Slide No. 15331, O69/1, H. *Caheniasaccites elongatus*, BSIP Slide No. 15330, E63/1, I. *Scheuringipollenites maximus*, BSIP Slide No. 15329, Q46/2, J. *S. maximus*, BSIP Slide No. 15328, K67/4, K. *S. maximus*, BSIP Slide No. 15325, K51, L. *Cycadopites* sp., BSIP Slide No. 15329, F52/4, M. *Striasulcites* sp., BSIP Slide No. 15329, F52/4, N. *Falcisporites zapfei*, BSIP Slide No. 15332, W59/2, O. *Parasaccites densicarpus*, BSIP Slide No. 15324, M61/4, P. *P. diffusus*, BSIP Slide No. 15321, E50/2, Q. *Plicatipollenites ovatus*, BSIP Slide No. 15332, W55/3, R. *P. indicus*, BSIP Slide No. 15334, P64/4.




Figure 5. Stratigraphically significant palynotaxa of Wordian-Changhsingian

A. *Striatopodocarpites multistriatus*, BSIP Slide No. 15325, G49/1, B. *S. multistriatus*, BSIP Slide No. 15325, M39/1, C. *Lunatisporites pellucidus*, BSIP Slide No. 15319,

S61/2, D. *Striatites rhombicus*, BSIP Slide No. 15325, C37/4, E. *Guttulapollenites hannonicus*, BSIP Slide No. 15325, O63/4, F. *Corisaccites alutus*, BSIP Slide No. 15325, O36/4, G. *C. alutus*, BSIP Slide No. 15334, R60/2, H. *Praecolpatites nidpurensis*, BSIP Slide No. 15321, O68/4, I. *Lunatisporites* sp., BSIP Slide No. 15331, N69/4, J. *Horriditriletes ramosus*, BSIP Slide No. 15331, H49/3, K. *Latosporites colliensis*, BSIP Slide No. 15319, V52/4, L. *Maculatasporites* sp., BSIP Slide No. 15335, J61/1, M. *Strotersporites perfectus*, BSIP Slide No. 15326, H41/2, N. Unidentified bisaccate-monolete, BSIP Slide No. 15333, G53/1, O. *Faunipollenites singrauliensis*, BSIP Slide No. 15326, F43/3, P. *Vestigipollenited rudis*, BSIP Slide No. 15326, Q36/2.

Palynological assemblage-I (Striate bisaccates- *Striasulcites* spp.)

The distance from 429.50 m to 232.10 m is a part of this palynoassemblage. It is distinguished by a predominance of striate bisaccate pollen grains, primarily species of *Faunipollenites* (= *Protohaploxypinus*) spp. (13-36%) {*F. bharadwajii*, *F. varius*, *F. enigmatus*, *F. singrauliensis*}, *Striatopodocarpites* spp. (1-10%) {*S. brevis*, *S. multistriatus*, *S. diffuses*, *S. magnificus*} and *Striasulcites* spp. (5-50%) {*S. tectus*, *S. ovatus*}. The other taxa of this assemblage that have been recorded are:

Triletes: *Latosporites colliensis* (1-3%), *Calamospora exile* (1%), *Lophotriletes* sp. (1%), *Horriditriletes* sp. (1%), *Lahirites* sp. (1%), *Leiotriletes* sp. (1%), *Microfoveolatispora foveolata* (1%).

Monosaccates: *Caheniasaccites elongatus* (1-3%), *Densipollenites* spp. (1-10%) {*D. densus*, *D. magnicarpus*, *D. invisus*, *D. marginalis*}, *Parasaccites* (1-7%) {*P. densicarpus*, *P. distinctus*, *P. korbaensis*, *P. diffusus*}, *Crucisaccites* sp. (2%), *Plicatipollenites* sp. (1%), *Potonieisporites* sp. (1%), *Striomonosaccites ovatus* (1-5%).

Non-striate bisaccates: *Klausipollenites* sp. (1-5%), *Alisporites* sp. (1-7%), *Chordasporites australiensis* (1-7%), *Falcisporites nuthaliensis* (1-4%), *Scheuringipollenites* spp. (2-28%)

{*S. barakarensis*, *S. tentulus*, *S. maximus*}, *Ibisporites diplosaccus* (1-7%), *Brachysaccus* sp. (1%), *Primuspollenites levis* (1%), *Vestigisporites rudis* (1%), *Platysaccus* sp. (1%).

Striate bisaccates: *Crescentipollenites globosus* (1-5%), *Strotersporites communis* (2-13%), *Striasulcites tectus* (2-3%), *Striatites rhombicus* (1%).

Taeniates: *Lunatisporites* spp. (1-4%) {*Lunatisporites* sp., *L. pellucidus*}, *Corisaccites alutus* (2%), *Guttulapollenites hannonicus* (1%), *Hamiapollenites insolitus* (1%), *Kamthisaccites kamthiensis* (1%).

Others: *Inaperturopollenites* sp. (1-7%), *Weylandites indicus*, *W. lucifer* (1%), *Cuneatisporites* sp. (1-3%), *Cycadopites* sp. (1%), *Praecolpatites nidpurensis* (1%).

Comparison: Palynological assemblage-I compares well with the *Gondisporites raniganjaensis* Zone of the Damodar Basin (Tiwari & Tripathi, 1992). The assemblage also has its resemblance with other records (Palynoassemblage-II: Srivastava and Jha, 1988; Palynozone-F: Tiwari and Ram-Awatar, 1989; Assemblage-IV: Tiwari *et al.*, 1991; Palynozone 2: Srivastava and Bhattacharyya, 1996; Palynozone-5: Jha and Aggarwal, 2012; Palynozone-6: Aggarwal and Jha, 2013; Palynoassemblage-III: Jha and Aggarwal, 2015; Assemblage-I: Mishra *et al.*, 2018; Palynoassemblage-6: Jha *et al.*, 2018; Palynoassemblage-IV: Aggarwal *et al.*, 2019).

Lithostratigraphic distribution: Kamthi Formation.

Age: *Gondisporites raniganjaensis* closely resembles Zone-VII of the Western Australian Gondwana palynological zones (Kemp *et al.*, 1977) in the predominance of the striate bisaccates *Faunipollenites* (= *Protohaploxypinus*) spp., *Striatites* spp. and *Striatopodocarpites* spp. and monosaccate *Densipollenites* spp. Zone-VII of Western Australia has been dated as Wordian-Capitanian (Guadalupian), and Zone-VI has been precisely dated as Roadian based on the recovered ammonites and other invertebrate fossils that lie just below this zone.

Wordian – Capitanian (Guadalupian) age has been thus assigned for palynoassemblages-I. Similar records have also been documented in Indian Gondwana (Table 5; Aggarwal and Jha, 2013, Fig. 2; Prasad and Pundir, 2017).

Interregional correlation: By sharing occurrences of *Striatopodocarpites* spp., *Faunipollenites* spp., *Densipollenites* spp., *Scheuringipollenites* spp., *Guttulapollenites hannonicus*, and *Weylandites lucifer*, the current Jangareddygudem palynoflora of India is tentatively correlatable with the Late Permian palynoassemblages of (Balme and Playford, 1967; Kemp, 1973; Dibner, 1976; Playford, 1990). It corresponds to the NW Nunatak, Vestfjella, and southern section palynological assemblages from the Late Permian in Dronning Maud Land, Antarctica (Lindström, 1996). *Striatopodocarpites* spp. and *Faunipollenites* spp., which are dominant in both assemblages, are supplemented by a few non-striate bisaccates (*Scheuringipollenites* spp., *Alisporites* spp., etc.). The palynoflora of the Buckley Formation (Mount Archenar, Central Transantarctic Mountains, Farabee *et al.*, 1991) and the top portions of the Mount *Glossopteris* Formation and Queen Maud Formation (Nilsen Plateau, Kyle, 1977; Kyle and Schopf, 1982) from the Late Permian are exhibiting notable parallels. The present palynoflora indicates similarities in having commonly occurring palynocomponents to some extent, with the Upper stage-5 of Australia represented by the dominance of striate bisaccates and a few non-striate bisaccates along with trilete spores (Foster, 1982). The latest Permian palynoassemblages of Australia (Backhouse, 1993) are documented to show *Horriditriletes* spp., *Protohaploxypinus* (= *Faunipollenites* spp.), and *Weylandites lucifer*, which forces for the broad tentative correlation with Indian palynoflora. Present palynoflora has also been correlated with the African palynoflora of Assemblage Zone IV (Falcon, 1975), with the upper part of the Madumabisa Formation palynoassemblage, Zambia (Utting, 1976), and microfloral zone 6 and 7 (Anderson, 1977). Further, the palynoflora of the lower Sakamena Group, Madagascar (Wright and Askin,

1987), and Biozone-E of the north Karoo Basin (MacRae, 1988), including the Palynoassemblage of the Lower Karoo sequence (Semkiwa *et al.*, 2003) and palynoassemblage of the Kalahari Karoo Basin (Modie and Le Hérisse, 2009) are depicting palynofloral similarities. It also indicates little correlation similarity with the *Tornopollenites toreutos* Zone, Andira' Formation of Tapajo's Group, Amazon Basin, northern Brazil (Playford and Dino, 2000, 2005).

Palynological assemblage-II (Striate bisaccates- *Densipollenites* spp.)

The Palynological assemblage-II (208.30-83.50m) is distinguished from the previous assemblage by the predominance of striate bisaccates, *Striatopodocarpites* spp. (2-24%) {*S. tiwarii*, *S. brevis*, *S. subcircularis*, *S. multistriatus*, *S. cancellatus*}, *Faunipollenites* (= *Protohaploxypinus*) spp. (13-40%) {*F. parvus*, *F. varius*, *F. goraiensis*, *F. bharadwajii*} along with a significant percentage of enveloping monosaccate *Densipollenites* spp. (6-33%) {*D. indicus*, *D. invisus*, *D. magnicarpus* var. *distinctus*, *D. magnicarpus* var. *annulatus*}. The composition of this palynoassemblage is as follows:

Triletes: *Latosporites* (1-23%), *Horriditriletes ramosus* (1%), *Lacinitriletes* sp., *Lophotriletes rectus* (1%), *Leiotriletes* sp. (1%), *Calamospora* sp. (1%).

Monosaccates: *Parasaccites* spp. (2%) {*P. obscures*, *P. korbaensis*}, *Plicatipollenites* spp. {*P. ovatus*, *P. indicus*}, *Potonieisporites* sp. (1%), *Striomonosaccites ovatus* (1%), *Crucisaccites* sp. (1%).

Non-striate bisaccates: *Scheuringipollenites* spp. (8-40%) {*S. barakarensis*, *S. tentulus*, *S. maximus*}, *Klausipollenites schaubergeri* (1-6%), *Alisporites* spp. (1-4%) {*A. landianus*, *A. indarraensis*}, *Chordasporites australiensis* (1-11%), *Falcisporites zapfei* (1-3%), *Ibisporites diplosaccus* (1-10%), *Platysaccus* sp. (1-3%), *Primuspollenites* sp. (1%), *Vestigisporites* sp. (1%), *Vesicaspora luteus* (1%).

Striate bisaccates: *Striapollenites saccates* (1-3%), *Striatites* spp. (1-5%) {*S. tentulus*, *S. parvus*, *S. communis*, *S. reticuloidus*}, *Striasulcites* spp. (1-4%) {*S. ovatus*, *S. tectus*}, *Crescentipollenites* spp. (3-11%) {*C. globosus*, *C. barakarensis*, *C. densus*}, *Strotersporites* spp. (1-19%) {*S. wilsonii*, *S. communis*, *S. crassiletus*, *S. perfectus*}, *Lahirites* sp. (1%), *Verticopollenites* spp. (2%) {*V. debilis*, *V. secretus*, *V. finimitus*}.

Taeniates: *Corisaccites alutus* (1%), *Hamiapollenites insolitus* (1%), *Lunatisporites ovatus* (1%), *Guttulapollenites* spp. (1%) {*G. hannonicus*, *G. gondwanensis*}, *Lueckisporites virkkiae* (1%).

Others: *Inaperturopollenites* sp. (1-5%), *Cuneatisporites* sp. (1-4%), *Weylandites circularis* (1%), *Rhizomaspora indica* (1%), *Staurosaccites* sp. (2-3%), *Tiwariasporis* sp. (1%), *Latosporites colliensis* (1%), *Maculatasporites* sp. (1%).

Lithostratigraphic distribution: Kamthi Formation.

Comparison: Palynoassemblage-II is akin to the *Densipollenites magnicarpus* Assemblage zone of Damodar Basin (Tiwari and Tripathi, 1991) in predominance of striate bisaccates along with acme of the monosaccate genus *Densipollenites* spp. It also closely resembles the *Guttulapollenites gondwanensis*-*Striatopodocarpites tiwarii* zone of Krishna Godavari Basin (Prasad *et al.*, 1995; Prasad and Pundir, 2017). Similar palynoflora has also been recorded from other places (Palynozone-6: Jha and Aggarwal, 2012; Palynozone-8: Srivastava and Jha, 1995; Palynoassemblage-II: Jha, 2004; Palynoassemblage-I: Jha *et al.*, 2014; Assemblage-II Mishra *et al.*, 2018; Palynoassemblage-VII: Jha *et al.*, 2018). Besides, this *Densipollenites* assemblage zone is well-known from the other Indian Gondwana basins such as South Rewa Basin (Ram-Awatar *et al.*, 2005); Damodar Basin (Srivastava *et al.*, 1997; Vijaya, 2011; Murthy, 2010; Murthy *et al.*, 2010; Murthy *et al.*, 2015); Satpura Basin (Bharadwaj *et al.*, 1978; Murthy *et al.*, 2013); Mahanadi Basin (Tiwari *et al.*, 1991; Tripathi, 1997; Chakraborty, 2003; Vijaya *et al.*, 2012; Sahoo *et al.*, 2020); Rajmahal Basin (Vijaya, 2006,

2009; Tripathi *et al.*, 2010); Talcher Basin (Patel *et al.*, 2022); Son Valley (Tiwari and Ram-Awatar, 1989).

Age: However, the *G. gondwanensis*- *S. tiwarii* zone documented from the latest Permian (uppermost part of Lower Gondwana succession) in India, including Godavari Coalfield, is also closely comparable with the Late Permian palynofloras of Chhidru Formation (Salt Range, Pakistan: Balme, 1970) and Gungri Shale (Spiti Valley, Tethyan Himalaya: Singh *et al.*, 1995) in having the dominant occurrence of the species of *Striatopodocarpites*, *Densipollenites*, *Crecentipollenites*, and *Guttulapollenites*. The associated sediments have been dated as Lopingian (Wuchiapingian-Changhsingian) based on the occurrence of the *Productus*, *Spiriferella*, and *Cyclolobus walkeri* (Prasad and Pundir, 2017). The occurrence of *Densipollenites magnicarpus* zone just below the *Lunatisporites pellucidus*–*Klausipollenites schaubergeri* Zone of Griesbachian (Early Triassic) age in India (= *Lunatisporites pellucidus* Zone; Helby *et al.*, 1987) confirms its Wuchiapingian-Changhsingian (Lopingian) age. So, the Lopingian age has been assigned to the palynoassemblage-II of this study (Table-5: Aggarwal and Jha, 2013, Fig. 2; Prasad and Pundir, 2017).

International correlation: Present palynoassemblage is tentatively correlated with the Late Permian palynoflora of the Prince Charles Mountain (Lindström & McLoughlin, 2007) as well as with the Buckley Formation, Central Transantarctic Mountain (Farabee *et al.*, 1991; Ram-Awatar *et al.*, 2014). It shows similarity with the palynoassemblage reported by Collinson *et al.* (2006) from the base of Buckley Formation at Graphite Peak. The palynocomponents in the present assemblage allow its tentative correlation with the palynoflora of the African Maji Ya Chumvi Formation (Hankel, 1992), upper Stage 5 of Foster (1982) in Eastern Australia, and with the lower part of the Sabina Sandstone (Backhouse, 1993), with Unit-VIII of the Grant Formation, with the Amb Formation (Balme, 1970), Canning Basin (Kyle, 1977), and with the Sardhai Formation (Jan *et al.*, 2009) along

with the Chhidru Formation (Hermann *et al.*, 2011, 2012, 2015) of the Salt Range. Correlation of the present palynoflora can be seen with the *Klausipollenites schaubergeri* Zone of Steiner *et al.* (2003) of Carlton Heights. It can be correlated with the assemblage described by Prevec *et al.* (2010) in New and Old Wapadsberg Pass, Eastern Cape Province, South Africa. It shows similar features compared with the Lower Sakamena Group palynoassemblage of Madagascar (Wright and Askin, 1987). However, Late Permian palynozones of South America have significant disparities with the Indian palynoflora. However, the present Late Permian Jangareddygudem palynoassemblage shows some similarities with the *Tornopollenites toreutos* Zone, northern Brazil (middle and upper part of the Andirá Formation, Playford and Dino, 2005), and the La Veteada Formation, Argentina (Gutiérrez *et al.*, 2011) in having a common occurrence of diverse association of striate bisaccates and a few spores. It can broadly be correlated with the palynoassemblage of the La Veteada Formation, Sierra De Narváez Catamarca Province, Argentina (Zavattieri *et al.*, 2008).

PALYNOFACIES ANALYSIS

In the present study, the palynofacies are exclusive of continental origin as the organic matter is entirely devoid of any type of marine element (dinoflagellate cysts, foraminiferal linings, marine algae, etc.). The palynofacies components are represented by the pollen-spore (SP), which includes spores, pollen grains, fungal spores; structured phytoclasts (ST)/translucent phytoclasts (wood particle, tracheids, cuticles, and planar organic matter); opaque black phytoclasts (charcoal) comprised of oxidized black to brownish-black coloured woody tissues, including tracheids; degraded organic matter (DOM) includes pale yellow to yellow/brown disintegrated organic matter produced after degradation of the algal/fungal activity, whose cellular structure is not clear; amorphous organic matter (AOM) includes all

microscopic structureless organic components including bacterially derived AOM, resins and amorphous products of the diagenesis of macrophytic tissues.

Samples have been processed for the palynofacies analysis, out of which 35 samples were found rich in organic matter, and two samples (6 & 17) were low in organic matter. The configuration of various organic components is plotted in a ternary diagram proposed by Hacquebard and Donaldson (1969) and modified by Singh and Singh (1996) and Aggarwal *et al.* (2019) to link the current outcomes to the depositional setting. The frequency distribution of organic matter in borehole MJR-13 has been given in Fig. 6. The different types of palynofacies components recovered from the studied sequence of the Jangareddygudem area of Godavari Coalfield are shown in Fig. 7. Based on the recovery of the different types of organic matter, four distinct (Fig. 8) palynofacies assemblages (PF-I to PF-IV) have been identified, which are described as follows.




Figure 6 Frequency distribution of organic matter and palynofacies identified in borehole MJR-13.




Figure 7. Representative photomicrographs of palynofacies components (all bars 50µm). A. Palynofacies showing palynomorphs and degraded terrestrial phytoclasts, BSIP Slide No. 15327, Q55/1, B. translucent wood, BSIP Slide No. 15329, K45/3, C. translucent wood, BSIP Slide No. 15328, L43/2, D. opaque wood phytoclasts, BSIP Slide No. 15329, J60/4, E. structured organic matter (leaf cuticle), BSIP Slide No. 15335, P63/2, F. Structured organic matter, BSIP Slide No. 15321, W49/1, G. degraded terrestrial matter, BSIP Slide No. 15333, M60/2, H. degraded terrestrial organic matter, BSIP Slide No. 15331, D45/2, I. Amorphous organic matter (AOM), BSIP Slide No. 15325, S41/1, J. Amorphous organic matter (AOM), BSIP Slide No. 15321, S45/1.

Palynofacies-I (PF-I)

This palynofacies is the most occurred (18 samples; Fig. 8) palynofacies assemblage in the whole succession, which is mainly predominated by the occurrence of the ST (3.78-79.07%; av. 33.84%), followed by the occurrence of the SP (0-49.79%; av. 26.37%). Other constituents of this assemblage are DOM (0-46.66%; av. 27.17%), opaque phytoclasts (0-37.80%; av. 10.67%) and AOM (0- 4.6%; av. 1.89%). Along with these elements, a meager percentage (0-0.87%; av. 0.71%) of fungal remains has also been documented. PF-I corresponds to the carbonaceous shales, shaly coals, and very fine to fine-grained sandstone.

Palynofacies-II (PF-II)

PF-II is featured by the predominance of the degraded matter, including DOM (1.7-69.56%; av. 41.32%) and AOM (0.3-53.81%; av. 12.32%) and sub-dominated by the occurrence of the SP (7.8-34.72%; av. 17.19%) and ST (av. 15.63%). Opaque phytoclasts of PF-II are in fewer amounts (0.8-24.62%; av. 13.50%). This type of association is represented by the shale, very fine to medium-grained sandstone. PF-II is represented by the five samples (Fig. 8).

Palynofacies-III (PF-III)

PF-III is shown by the abundant occurrence of opaque phytoclasts (62.70-97.34%; av. 82.02%). Other constituents of this palynofacies association are ST (2.53-14.73%; av. 9.10%), DOM (0.8-12.29%; av. 4.74%), AOM (0-7.25%; av. 2.19%) and SP (0-4.9%; av. 1.93%). PF-III is mainly represented by coarse-grained sediments like coarse-grained sandstone, and sandstone with carbonaceous matter except in one or two places (shale and thin coal band). A total of eight samples of this succession are represented by PF-III (Fig. 8).

Palynofacies-IV (PF-IV)

PF-IV is the least occurred palynofacies in borehole MJR-13. This palynofacies association is represented by the predominance of the opaque phytoclasts (43.23-47.06%; av. 45.28%) sub-dominated by ST (15.72-45.09%; av. 36.20%). Other constituents of PF-IV are SP (0-15.72%; av. 4.98%), DOM (5.88-23.58%; av. 11.60%), and AOM (0-5.91%; av. 1.91%). This type of association is represented by various lithologies like claystone, shale, and fine to medium-grained sandstone (Fig. 8).




Figure 8. Ternary diagram showing the distribution of various palynofacies (after Aggarwal et al., 2019).

PALAEOENVIRONMENTAL INTERPRETATION

The palynofloral analysis demonstrates the predominance of the glossopterid pollen grains and the abridged proportion of pteridophytic spores. The gymnospermous pollen grains are represented mainly by monosaccates, monosulcate, striate bisaccates, non-striate bisaccates, taeniates, praecolpate and costate grains. The abundant occurrence of the arborescent vegetation suggests the hygrophilous to the mesophilous environment (Cazzulo-Klepzig *et al.*, 2005; Ruckwied *et al.*, 2014) in the recovered palynofloral assemblages (Fig. 2 and 3). Predominant occurrence of the gymnospermous pollen grains and the scarce presence of the algal (*Inaperturopollenites* sp., *Maculatasporites* sp. and *Latosporites colliensis*) and

peridophytic (*Calamospora*, *Horriditriletes*, *Lacinitriletes*, *Leiotriletes*, *Lophotriletes*, *Microfoveolatispora*) forms also suggests the hygrophilous to mesophilous environmental (Rothwell, 1988; Di Michele and Philips, 1994) conditions for both of the palynoassemblages (Fig.9, 10 and Table-2).




Figure 9. The number of palynotaxa recovered in borehole MJR-13 belongs to different plant groups.




Figure 10. Differential distribution of palynotaxa belonging to different plant groups at individual depth levels.

Interpretation of the depositional environment based on palynofacies analysis:

Palynofacies assemblage-I is characterized by the dominant occurrence of ST and SP. By taxonomic characterization, palynomorphs can usually be identified and assigned their botanical affinities, while structured phytoclast particles with internal structures can only be related to a type of plant. The dominance of gymnospermous palynomorphs and structured organic matter (dominated by the wood particles and tracheids, Fig. 8) having an affinity with sub-arborescent/arborescent vegetation suggests proliferated forest development with its closeness to the terrestrial landmass (due to the absence of any marine element) in the temperate climatic conditions (Götz and Ruckwied, 2014).

The dominance of structured organic matter derived from higher plant debris implies a fairly dense vegetation cover near the deposition site. Land plant-derived phytoclasts are represented by structured and opaque phytoclasts, which may be equidimensional or elongated lath-shaped and translucent fragments of thin, tubular, elongated wood tracheids. Hydrodynamic equivalence of translucent wood phytoclasts controls their distribution in sediments since woody phytoclasts are composed of relatively larger and denser particles, and their abundances have commonly been reported to correlate with coarse silts and very fine-grain sand (Habib, 1983; Firth, 1993; Tyson, 1993). Their high relative and absolute abundances indicate a substantial influx with proximal depositional settings (Muller, 1959; Pocklington and Leonard, 1979). The concurrence of sediments comprising carbonaceous shale, shale, fine-grained sandstone, and organic matter components has led to assuming low-energy environmental settings through the deposition of PF-I. Thus, overall, the swampy environments have been deduced for PF-I.

PF-II is represented by the dominance of degraded organic matter, including DOM and AOM, followed by SP and ST. Degraded plant debris and amorphous organic matter derived from the microbiological activity, indicating the existence of dysoxic-anoxic conditions

during the deposition of sediments. AOM is characterized by a lack of cellular structures and indistinct and unrecognizable outlines. Structureless organic matter cannot be inferred from its probable botanical affinity and has also been documented in terrestrial settings (Tyson, 1995). High relative or absolute abundances of DOM and AOM are usually associated with deposits below the upwelling water column and dysoxic conditions in the bottom (Davey and Rogers, 1975; Tissot and Pelet, 1981; Summerhayes, 1983). AOM has been reported to decrease in shallow shelf sediments, whereas it tends to increase in a deeper basinward direction under dysoxic-anoxic conditions (Dow and Pearson, 1975; Bujak *et al.*, 1977). According to Batten and Stead (2002), in freshwater and brackish environments, oxidizing and high or low energy conditions may be associated with periodically exposed areas of floodplains, riverbeds, and lakes or lagoon margins (Zhang *et al.*, 2010). The concurrence of the sedimentological data and organic matter components in high to low energy conditions in the flooded palaeomires have been inferred for PF-II.

PF-III is dominated by opaque phytoclasts. The hydrodynamic equivalence of opaque woody phytoclasts has been indicated to control by particle size rather than its shape (Tyson, 1995). Larger particles have been reported to be collected in proximal and relatively high-energy environments, i.e., sand and silt (Van der Zwan, 1990; Baird, 1992). Opaque wood phytoclast concentrations have been found to reflect deposition polarity, sediment transport distance, and oxygenation level in host sediments (Carvalho *et al.*, 2013). According to numerous studies, (including those by Marsan and Pocock, 1981; Pocock *et al.*, 1987; Closas *et al.*, 2005; and Piekowski and Waksmundzka, 2009), opaque phytoclasts are frequently found in oxic swamps, prodelta, delta fronts, and river deposits. Black wood particles are very common in high-energy, proximal, coarse-grained sediments of fluvial to delta-top systems (Fisher, 1980; Nagy *et al.*, 1984; Smyth *et al.*, 1992; Williams, 1992). Except one or two locations (Fig. 2), PF-III was largely dominated by coarser sediments, which may have been

caused by the abundance of terrigenous clastic deposits due to increased fluvial input from precipitation. The considerable representation of opaque phytoclasts and the low occurrence of other organic matter types in PF-III suggest that exposed sections of deposited beds experienced extremely oxidizing conditions. Most of the lath-shaped opaque phytoclasts over the equidimensional ones have been noticed in the present studies, reflecting deposition in proximal settings (Batten and Stead, 2002). Oxidized swamp settings have been thus inferred for this palynofacies assemblage (Fig. 8).

The sub-dominance of the ST and the dominance of the OP are characteristics of PF-IV. The predominance of OP in PF-IV demonstrates its deposition in oxidizing environments, while the sub-dominance of ST depicts the deposition of this form of palynofacies association close to the source of the vegetation. As a result, it was concluded that PF-IV was deposited under less oxidizing conditions than PF-III. The concurrence of the lithological data and organic matter elements for PF-IV demonstrates the mixed environmental setup, which mainly prevailed by the oxic swamps with low to medium energy levels.

Palynofacies study results demonstrate that the succession of MJR-13 is dominated by the presence of PF-I (Fig. 8), which clearly demonstrates swampy settings in a dysoxic to anoxic environment for the succession and corroborates earlier research conducted by investigators during this time (Aggarwal *et al.*, 2019).

CONCLUSIONS

1. Based on the palynofloral evidence, two distinct palynoassemblages (I-II) have been identified within the Kamthi Formation located in the Chintalapudi sub-basin of the Godavari Coalfield. Palynoassemblage-I (*Striates*+*Striasulcites* spp.), and Palynoassemblage-II (*striates*+*Densipollenites*), which have been dated as Guadalupian and Lopingian, respectively.

2. Based on the recovered different types of organic matter, four distinct palynofacies (I-IV) have been identified. These palynofacies are characterized by the dominance of structured phytoclasts along with sporomorphs; degraded OM along with amorphous OM; opaque phytoclasts and opaque phytoclasts along with structured phytoclasts, respectively.
3. The presence of freshwater algal spores and the complete absence of marine components in the palynological samples attested to the deposition in the freshwater regime.
4. Palynofacies-I indicates dense vegetation cover close to the site of deposition with its closeness to the terrestrial landmass in swampy settings; PF-II suggests the existence of dysoxic-anoxic conditions in the flooded palaeomires; PF-III indicates oxidized swamp settings, and PF-IV represents the mixed environmental setup which is mainly prevailed by the oxic swamps with low to medium energy levels.
5. Succession is mainly predominated by the abundant occurrence of the glossopterids and conifers co-occurring along with filicophytes and sphenophytes is indicative of hypoautochthonous sedimentation in the proximal settings and low-energy environments. It denotes that the landmass was located in more inland areas, and peat formation occurred in forest swamps in the terrestrial regime.

Acknowledgements: *Authors express their sincere gratitude to the Director, Birbal Sahni Institute of Palaeosciences, for extending the necessary research facilities used during the studies. Sri M. Basva Chari, former Chief General Manager, Singareni Collieries Company Limited, is thankfully acknowledged for his cooperation, fruitful discussion, and facilities extended during field visits and sample collection in various areas providing the informative details of geology and location of drilled boreholes, etc. The members of the Exploration Division are also acknowledged for their cooperation during the field visit. Sri ADP Rao, Mineral Exploration Corporation Limited, rendered generous help during the sample*

collection and acknowledged thankfully for providing necessary field details. The authors humbly express gratitude to the anonymous reviewers for dedicating their valuable time and providing insightful comments that have significantly enhanced the quality and content of this manuscript.

REFERENCES

Aggarwal N & Jha N 2013. Permian palynostratigraphy and palaeoclimate of Lingala Koyagudem Coalbelt, Godavari Graben, Andhra Pradesh, India. *Journal of Asian Earth Sciences* 64: 38-57.

Aggarwal N, Agrawal S & Thakur B 2019. Palynofloral, palynofacies and carbon isotope of Permian coal deposits from the Godavari Valley Coalfield, South India: Insights into the age, palaeovegetation and palaeoclimate. *International Journal of Coal Geology* 214: 1-22.

Anderson JM 1977. The biostratigraphy of the Permian and Triassic Part 3. A review of Gondwana palynology with particular reference to the northern Karoo Basin, South Africa. *Memoir Botanical Survey of South Africa* 41: 1-188.

Backhouse J 1993. Palynology and correlation of Permian sediments in the Perth Collie, and Officer Basins, Western Australia. *Report Geological Survey of Western Australia* 34: 111-128.

Baird JG 1992. Palynofacies of the eastern margin of Gippsland Basin-Energy, Economics, and Environment. In: *Proceeding of the Gippsland Basin*, Melbourne June, Australian Institute of Mining and Metallurgy 25-42.

Balme BE 1970. Palynology of Permian and Triassic strata in the Salt Range and Surghar Range, west Pakistan; In Kummel, B. & Teichert, C. (eds.): "Stratigraphic boundary

problems: Permian and Triassic of West Pakistan", Department of Geology, University of Kansas. Special publication 4: 305-453.

Balme BE 1995. Fossil in situ spores and pollen grains: An annotated catalogue. Review of Palaeobotany and Palynology 87: 81–323.

Balme BE & Playford G 1967. Late Permian plant microfossils from the Prince Charles Mountains, Antarctica. Revue de Micropaléontologie 10: 179–192.

Batten DJ 1996. Palynofacies and palaeoenvironmental interpretation-26 A. In Jansonius, J. & McGregor, D.C. (eds.): Palynology-Principles and applications. American Association of Stratigraphic Palynologists Foundation 3: 1011–1064.

Batten DJ 1999. Palynofacies analysis, In Jones, T.P. & Rowe, N.P. (eds.): Fossil plants and Spores: Modern Techniques. London Geological Society 194-198.

Batten DJ & Stead DT 2002. Palynofacies analysis and its stratigraphic application; In Koutsoukos, E.A.M. (ed.): Springer, Dordrecht, Applied Stratigraphy 203–226.

Bharadwaj DC 1962. The miospore genera in the coals of Raniganj Stage (Upper Permian) India. The Palaeobotanist 9 (1-2): 68-106. <https://doi.org/10.54991/jop.1960.600>

Bharadwaj DC & Salujha SK 1964. Sporological study of seam VIII in Raniganj Coalfield, Bihar (India) Part I- Description of sporae dispersae. The Palaeobotanist 12 (2): 181-215. <https://doi.org/10.54991/jop.1963.653>

Bharadwaj DC, Tiwari RS & Anand-Prakash 1978. Palynology of Bijori Formation (Upper Permian) in Satpura Gondwana Basin, India. The Palaeobotanist 25: 70-78. <https://doi.org/10.54991/jop.1976.997>

Birks HJB & Birks HH 1980. Quaternary Palaeoecology-2a. In Arnold, E. (ed.): London Press 289.

Blanford WT 1872. A note on plant-bearing sandstones of the Godavari on the southern extension of rocks belonging to Kamthi Group to the neighborhood of Eluru and Rajahmundry on the possible occurrence of coal in the same directions. Record Geological Survey of India 4: 59-88.

Bujak JP, Barss MS & Williams GL 1977. Offshore east Canada's organic type and color and hydrocarbon potential. The Oil and Gas Journal 75: 198-201.

Carvalho Marcelo de A, Ramos RRC, Crud MB, Witovisk L, Kellner AWA, Silva Helder de P, Grillo ON, Riff D & Romano PSR 2013. Palynofacies as indicators of paleoenvironmental changes in a Cretaceous succession from the Larsen Basin, James Ross Island, Antarctica. Sedimentary Geology 295: 53–66.

Cazzulo-Klepzig M, Menegat E & Guerra-Sommer M 2005. Palynology and palaeobotany in the reconstruction of landscape units from the candiota coalfield, Permian of Parana Basin, Brazil. Revista Brasileira de Paleontologia 8 (1): 83–98.

Chakraborty, S., 2003. Miofloral assemblage of the subsurface Lower Gondwana rocks of Permian in parts of Mahanadi Basin, India. Acta Palaeontologica Sinica 42, 13–21.

Closas MC, Permanyer A & Vila MJ 2005. Palynofacies distribution in a lacustrine basin. Geobios 38: 197–210.

Collinson JW, Hammer WR, Askin RA & Elliot DH 2006. Permian–Triassic boundary in the Central Transantarctic Mountains, Antarctica. Bulletin Geological Society of America 118: 747–763.

Costamagna LG, Kustatscher E, Scanu GG, Scanu MD, Rio P, Pittau P & van Konijnenburg-van Cittert JHA 2018. A palaeoenvironmental reconstruction of the Middle Jurassic of Sardinia (Italy) based on integrated palaeobotanical, palynological and lithofacies data assessment. *Palaeobiodiversity and Palaeoenvironments* 98: 111-138.

Davey RJ & Rogers J 1975. Palynomorph distribution in recent offshore sediments along two traverses of South West Africa. *Marine Geology* 18: 213–225.

Di Michele WA & Philips TL 1994. Palaeobotanical and palaeoecological constraints on models of peat-formation in the Late Carboniferous in Euramerica. *Palaeogeography Palaeoclimatology Palaeoecology* 106: 39–90.

Dibner AF 1976. Late Permian palynofloras in sediments around Beaver Lake (East Antarctica). *Antarctica* 15: 41–52.

Dow WG Pearson DB 1975. Organic matter in Gulf coast sediments; Reprints, Seventh Annual Offshore Technology Conference. *Offshore Technology Conference* 23-43.

Farabee MJ, Taylor EL & Taylor TN 1991. Late Permian palynomorphs from the Buckley Formation, Central Transantarctic Mountains, Antarctica. *Review of Palaeobotany and Palynology* 69: 353–368.

Falcon RMS 1975. Palynostratigraphy of the Karoo sequence in the central Sebungwe District, Mid Zambezi Basin. *Palaeontologica Africana* 18: 1–29.

Firth JV 1993. Palynofacies and thermal maturation analysis of sediments from the Nankai Trough; *In* Hill, I.A., Taira, A., Firth, J.V. et al. (eds.): *Proceedings of the Ocean Drilling Project, Scientific Results College Station, TX (Ocean Drilling Program)* 131: 57–69.

Fisher MJ 1980. Kerogen distribution and depositional environments in the Middle Jurassic of Yorkshire UK. *In* Singh, B.D., Singh, H.P. & Tiwari, R.S. (eds.): 4th International Palynological Conference, Lucknow 1976-1977.

Foster CB 1982. Spore-pollen assemblages of the Bowen Basin, Queensland (Australia): Their relationship to the Permian/Triassic boundary. *Review of Palaeobotany and Palynology* 36: 165–183.

Götz AE & Ruckwied K 2014. Palynological records of the early Permian postglacialclimate amelioration (Karoo Basin, South Africa). *Paleobiodiversity and Paleoenvironment* 94 (2): 229–235.

Gutiérrez P, Zavattieri AM, Ezpeleta M & Astini RA 2011: Palynology of the LaVeteda Formation (Permian) in the Sierra de Narvárez, Catamarca Province, Argentina. *Ameghiana* 48: 154–176.

Habib D 1983: Sedimentation-Rate-Dependent Distribution of Organic-Matter in the North-Atlantic Jurassic-Cretaceous. *Initial Reports of the Deep-Sea Drilling Project* 76: 781–794.

Hacquebard PA & Donaldson JR 1969. Carboniferous coal deposition associated with floodplain and limnic environments in Nova Scotia. *In*: Dapples, E.C. & Hopkins, M.E. (eds.): *Environment of Coal Deposition*. Geological Society of America Special Papers 114: 143–191.

Hankel O 1992: Late Permian to Early Triassic microfloral assemblages from the Maji Ya Chumvi Formation, Kenya. *Review of Palaeobotany and Palynology* 72: 129–147.

Helby R, Morgan R & Partridge AD 1987. A palynological zonation of the Australian Mesozoic. *Memoir of the Association of Australasian Paleontologists* 4: 1–94.

Hermann E, Hochuli PA, Bucher H & Roohi G 2012. Uppermost Permian to Middle Triassic palynology of the Salt Range and Surghar Range, Pakistan. *Review of Palaeobotany and Palynology* 169: 61–95.

Hermann E, Hochuli PA, Méhay S, Bucher H, Brühwiler T, Ware D, Hautmann M, Roohi G, Rehman K & Yaseen A 2011. Organic matter and palaeoenvironmental signals during the Early Triassic biotic recovery: the Salt Range and Surghar Range records. *Sedimentary Geology* 234: 19–41.

Hermann E, Kürschner WM, Kerp H, Bomfleur B, Peter A, Hochuli PA, Bucher H, Waree D & Ghazala R 2015. Vegetation history across the Permian-Triassic boundary in Pakistan (Amb section, Salt Range). *Gondwana Research* 27: 911–924.

Jan IU, Stephenson MH & Khan FR 2009. Palynostratigraphic correlation of the Sardhai Formation (Permian) of Pakistan. *Review of Palaeobotany and Palynology* 158: 72–82.

Jha N & Aggarwal N 2015. Peat-forming environment of coal-bearing Permian sediments in Kachinapalli area of Godavari Graben, India. *Revista Brasileira de Paleontologia* 18 (2): 239–250.

Jha N 2002. Palynological dating of sediments from Gattugudem area, Chintalapudi Sub-basin, Andhra Pradesh. *Geophytology* 30: 85–89.

Jha N 2004. Palynodating of coal-bearing sediments from the Bottapagudem area, Chintalapudi Sub-basin, Andhra Pradesh. *The Palaeobotanist* 53: 61–67.
<https://doi.org/10.54991/jop.2004.210>

Jha N 2008. Permian-Triassic palynofloral transition in the Sattupalli area, Chintalapudi Sub-basin, Godavari Graben, Andhra Pradesh, India. *Journal of Palaeontological Society of India* 52: 159–168.

Jha N & Aggarwal N 2012. Permian-Triassic palynostratigraphy in Mailaram area, Godavari Graben, Andhra Pradesh, India. *Journal of Earth System Science* 121: 1257–1285.

Jha N, Aggarwal N & Mishra S 2018. A review of the palynostratigraphy of Gondwana sediments from the Godavari Graben, India: Global comparison and correlation of the Permian-Triassic palynoflora. *Journal of Asian Earth Sciences* 163: 1-21.

Jha N, Pauline SK, Aggarwal N & Mahesh S 2014. Late Permian palynology and depositional environment of the Chintalapudi Sub-basin of Godavari Basin, Andhra-Pradesh, India. *Journal of Asian Earth Sciences* 79: 382-399.

Kemp EM 1973. Permian flora from the Beaver Lake area, Antarctica. 1. Palynological examination of samples. Bureau of Mineral Resources, Geology and Geophysics (Australia) Bulletin 126: 7–12.

Kemp EM, Balme BE, Helby RJ, Kyle RA, Playford G & Price PL 1977. Carboniferous and Permian palynostratigraphy in Australia and Antarctica: a review. Bureau of Mineral Resources, Geology and Geophysics (Australia) Bulletin 2: 177–208.

Kyle RA 1977. Palynostratigraphy of the Victoria Group, South Victoria Land, Antarctica. *New Zealand Journal of Geology and Geophysics* 20: 1081–1102.

Kyle RA & Schopf JM 1982. Permian and Triassic palynostratigraphy of the Victoria group, Trans-Antarctic Mountains. In Craddock (ed.), *Antarctic Geoscience*. University Wisconsin Press, Madison 649–659.

Lakshminarayana G 1996. Stratigraphy and structural framework of the Gondwana sediments in the Pranhita-Godavari Valley, Andhra Pradesh. In Guha, P.K.S., Sengupta, S., Ayyasami, K. & Ghosh R.N. (eds.): *Ninth International Gondwana Symposium, Hyderabad*. Geological Survey of India. Oxford and IBH Publishing Co., New Delhi-Calcutta 311–330.

Lakshminarayana G & Murti KS 1990. Stratigraphy of the Gondwana Formation in the Chintalapudi Sub-basins, Godavari Valley, Andhra Pradesh, India. *Journal of Geological Society of India* 36: 13–25.

Lindström S 1996. Late Permian palynology of Fossil ryggen, Vestfjella, Dronning Maud Land, Antarctica. *Palynology* 20: 15–48.

Lindström S & McLoughlin 2007. Synchronous palynofloristic extinction and recovery after the end-Permian event in the Prince Charles Mountains, Antarctica: Implications for palynofloristic turnover across Gondwana. *Review of Paleobotany and Palynology* 145: 89–122.

Looy CV, Brugman WA, Dilcher DL & Visscher H 1999. The delayed resurgence of equatorial forests after the Permian–Triassic ecologic crisis. *Proceedings of the National Academy of Sciences* 96: 13857–13862.

MacRae CS 1988. Palynostratigraphical correlation between the Lower Karoo sequence of the Waterberg and Pafuri coal basins and the Hammanskraal plant macrofossil locality, Republic of South Africa. *Memoir Geological Survey of South Africa* 75: 1–217.

Marsan T & Pocock SAJ 1981. In: Brooks, J. (ed.), *The Classification of Plant-Derived Particulate Organic Matter in Sedimentary Rocks-Review in Organic Maturation Studies and Fossil Fuel Exploration*. Academic Press, London 145–176.

Mishra S Aggarwal N & Jha N 2018. Palaeoenvironmental change across the Permian-Triassic boundary inferred from palynomorph assemblages (Godavari Graben, south India). *Palaeobiodiversity and Palaeoenvironment* 98: 177–204. <https://doi.org/10.1007/s12549-017-0302-3>. Mishra S & Singh VP 2018. Palynology, palynofacies, and taphonomical studies of

Kamthi Formation, (Godavari Graben), southern India: Implications to biostratigraphy, palaeoecology, and depositional environment. *International Journal of Coal Geology* 195: 102-124. <https://doi.org/10.1016/j.coal.2018.05.011>.

Modie BN & Le Hérissé A 2009. Late Palaeozoic palynomorph assemblages from the Karoo Supergroup and their potential for biostratigraphic correlation, Kalahari Karoo Basin, Botswana. *Bulletin of Geosciences* 84 (2): 337–358.

Mukhopadhyay GMS, Roychowdhury M & Parui PK 2010. Stratigraphic correlation between different Gondwana Basins of India. *Journal Geological Society of India* 76: 251–266.

Muller J 1959. Palynology of Recent Orinoco Delta and shelf sediments: reports of the Orinoco Shelf expedition. *Micropaleontology* 5: 1–2.

Murthy S 2010. Palynostratigraphy of the Permian succession in Borehole RJS-2, Raniganj Coalfield, Damodar Basin, West Bengal. *Journal of Indian Geological Congress* 2: 83–90.

Murthy S, Chakraborti B & Roy MD 2010. Palynodating of subsurface sediments, Raniganj Coalfield, Damodar Basin, West Bengal. *Journal of Earth System Science* 119: 701–710.

Murthy S, Kavali PS & Bernardes-de-Oliveira MEC 2015. Latest Permian palynomorphs from Jharia Coalfield, Damodar Basin, India and their potential for biostratigraphic correlation. *Revue de Micropaléontologie* 58 (3): 167–184.

Murthy S, Vijaya & Vethanayagam SM 2013. Palynostratigraphy of Permian succession in the Pench Valley Coalfield, Satpura Basin, Madhya Pradesh, India. *Journal of Palaeontological Society of India* 58: 241–250.

Nagy J, Dypvic H & Bajaerke T 1984. Sedimentological and paleontological analyses of Jurassic North Sea deposits from deltaic environments. *Journal of Petroleum Geology* 7: 169–188.

Patel R, Goswami S, Aggarwal N. & Mathews RP 2022. Lower Gondwana megaflora, palynoflora and biomarkers from Jagannath Colliery, Talcher Basin, Odisha, India and its biostratigraphic significance. *Geological Journal* 57 (3): 986-1004. DOI: 10.1002/gj.4318.

Pieńkowski G & Waksmundzka M 2009. Palynofacies in lower Jurassic epicontinental deposits of Poland: tool to interpret sedimentary environments. *Episodes* 32: 21–32.

Playford G 1990. Proterozoic and Palaeozoic palynology of Antarctica: a review. *In* Taylor, T.N. & Taylor, E.L. (eds.): *Antarctic Palaeobiology: Its Role in Reconstruction of Gondwana*. Springer-Verlag, New York 51–70.

Playford G & Dino R 2000. Palynostratigraphy of Upper Palaeozoic strata (Tapajos Group), Amazonas Basin, Brazil: Part- II. *Palaeontographica Abt B* 255: 87–145.

Playford G & Dino R 2005. Carboniferous and Permian Palynostratigraphy; *In*: Koutsoukos, E. A. M. (ed.), *Springer, Dordrecht. Applied Stratigraphy* 23: 101–121.

Pocklington R & Leonard JD 1979. Terrigenous organic matter in sediments of the St. Lawrence Estuary and the Saguenay Fjord. *Journal of the Fisheries Research Board of Canada* 36: 1250–1255.

Pocock AAJ, Vasanthy G & Venkatachala BS 1987. Introduction to the study of particulate organic materials and ecological perspectives. *Journal of Palynology* 23–24: 167–188.

Prasad B, Jain AK & Mathur YK 1995. A standard Palynozonation scheme for the Cretaceous and pre-Cretaceous subsurface sediments of Krishna-Godavari Basin, India. *Geoscience Journal* 16 (2): 155–233.

Prasad B & Pundir BS 2017. Gondwana biostratigraphy of the Purnea Basin (Eastern Bihar, India), and its correlation with Rajmahal and Bengal Gondwana Basins. *Journal of the Geological Society of India* 90: 405–427.

Prasad V, Singh IB, Bajpai S, Garg R, Thakur B, Singh A, Saravanan N & Kapur VV 2013. Palynofacies and sedimentology-based high-resolution sequence stratigraphy of the lignite-bearing muddy coastal deposits (early Eocene) in the Vastan Lignite Mine, Gulf of Cambay, India. *Facies* 59 (4): 737–761.

Prevec R, Gastaldo RA, Neveling J, Reid SB & Looy CV 2010. An autochthonous glossopterid flora with latest Permian palynomorphs and its depositional setting in the Dicynodon Assemblage Zone of the southern Karoo Basin, South Africa. *Palaeogeography Palaeoclimatology Palaeoecology* 292: 391–408.

Qureshy MN, Brahmam NK, Gadse SC & Mathur BK 1968. Gravity analysis and the Godavari rift, India. *Bulletin of Geological Society of America* 79: 1221–1230.

Raiverman V, Rao MR & Pal D 1985. Stratigraphy and structure of the Pranhita-Godavari Graben. *Petroleum Asia Journal* 8: 174–189.

Raja Rao CS 1982. Coalfield of India-2. Coal resources of Tamil Nadu, Andhra Pradesh, Orissa and Maharashtra. *Bulletin Geological Survey of India Series A* 45: 9–40.

Ram Awatar, Tewari R, Agnihotri D, Chatterjee S, Pillai SK & Meena KL 2014. Late Permian and Triassic palynomorphs from the Allan Hills, central Transantarctic Mountains, South Victoria Land, Antarctica. *Current Science* 106: 988–996.

Ram-Awatar, Kumar M & Prakash N 2005. Palynological analysis of Lower Gondwana sediments exposed along the Umrer River South Rewa Basin, Madhya Pradesh, India. *The Palaeobotanist* 54: 87–97. <https://doi.org/10.54991/jop.2005.71>

Rothwell GW 1988. Cordaitales. *In* Beck, C.B. (ed.): Origin and evolution of Gymnosperms. Columbia University Press 273–297.

Ruckwied K, Götz AE & Jones P 2014. Palynological records of the Permian eccla Group (South Africa): Utilizing climatic icehouse–greenhouse signals for cross-basin correlations. *Palaeogeography Palaeoclimatology Palaeoecology* 413: 167–172.

Sahoo M, Aggarwal N & Goswami S 2020. Palynological investigation of the lower Gondwana outcrop near Gopalprasad, Odisha, India: an inference on the age, palaeovegetation and palaeoclimate. *Journal of Palaeontological Society of India* 65(1): 27–35.

Semkiwa P, Kalkreuth W, Utting J, Mpanju F & Hagemann H 2003. The geology, Petrology, palynology and geochemistry of Permian coal basins in Tanzania: 2. Sonwe-Kiwira Coalfield. *International Journal of Coal Geology* 55: 157–186.

Singh MP & Singh PK 1996. Petrographic characterization and evolution of the Permian coal deposits of the Rajmahal basin, Bihar, India. *International Journal of Coal Geology* 29: 93–118.

Singh T, Tiwari RS & Ram-Awatar 1995. Stratigraphy and palynology of Carboniferous-Permian-Triassic succession in Spiti valley, Tethys Himalaya, India. *Journal Palaeontological Society of India* 40: 55–76.

Smyth M, Jian FX & Ward CR 1992. Potential petroleum source rocks in Triassic lacustrine-delta sediments of the Gunnedah Basin, Eastern Australia. *Journal of Petroleum Geology* 15: 435–450.

Srivastava SC & Bhattacharyya AP 1996. Permian-Triassic palynofloral succession in subsurface from Bazargaon, Nagpur District, Maharashtra. *The Palaeobotanist* 43: 10–15.
<https://doi.org/10.54991/jop.1994.1172>

Srivastava SC & Jha N 1988. A Lower Triassic palynoassemblages from Budharam area, Godavari Graben, Andhra Pradesh, India. *Geophytology* 18: 124–125.

Srivastava SC & Jha N 1992. Palynostratigraphy of Permian sediments in Manuguru area, Godavari Graben, Andhra Pradesh, India. *Geophytology* 22: 103–110.

Srivastava SC & Jha N 1993. Palynostratigraphy of Lower Gondwana sediments in Cintelapudi Sub-basin, Godavari Graben, Andhra Pradesh. *Geophytology* 23: 93–98.

Srivastava SC & Jha N 1994. Palynological dating of Lower Gondwana sediments in Sattupalli area, Chintalapudi Sub-basin, Andhra Pradesh, India. *The Palaeobotanist* 42: 169–173. <https://doi.org/10.54991/jop.1993.1143>

Srivastava SC & Jha N 1995. Palynostratigraphy and correlation of Permian-Triassic sediments in Budharam Area, Godavari Graben, India. *Journal of Geological Society of India* 46: 647–653.

Srivastava SC, Anand-Prakash & Kar R 1997. Palynology of Permian-Triassic sequence in Iria Nala, Tatapani-Ramkola Coalfield, India. *The Palaeobotanist* 46: 75–80.
<https://doi.org/10.54991/jop.1997.1318>

Steiner MB, Eshet Y, Rampino MR & Schwindt DM 2003. Fungal abundance spike and the Permian–Triassic boundary in the Karoo Supergroup (South Africa). *Palaeogeography Palaeoclimatology Palaeoecology* 194: 405–414.

Summerhayes CP 1983. Sedimentation of organic matter in upwelling regimes; In: Thiede, J. & Suess, E. (eds.), *Coastal Upwelling: Its Sediments Record Part B: Sedimentary Records of Ancient Coastal Upwelling* New York, Plenum Press 29–72.

Tissot B & Pelet R 1981. Sources and fate of organic matter in ancient sediments; 26th *Congres International de Geologie, Colloque Geologie des Oceans*, Paris 1980. *Oceanologica Acta*, Special Issue, Actes 97–103.

Tiwari RS & Ram-Awatar 1989. Spores dispersal and correlation of Gondwana sediments in Jharkhand Coalfield, Son Valley Graben, Madhya Pradesh. *The Palaeobotanist* 37: 94–114.
<https://doi.org/10.54991/jop.1988.1604>

Tiwari RS & Tripathi A 1991. Marker assemblage zones of spore and pollen species through Gondwana Palaeozoic-Mesozoic sequence in India. *The Palaeobotanist* 40: 194–236.
<https://doi.org/10.54991/jop.1991.1773>

Tiwari RS, Tripathi A & Jana BN 1991. Palynological evidence for Upper Permian Raniganj Coals in western part of Talcher Coalfield, Orissa, India. *Current Science* 61: 407–420.

Tripathi A 1997. Palynostratigraphy and palynofacies analysis of subsurface Permian sediments in Talcher Coalfield, Orissa. *The Palaeobotanist* 46: 79–88.
<https://doi.org/10.54991/jop.1997.1351>

Tripathi A, Murthy S & Singh R 2010. Palynodating of coal-bearing strata near Kunda Pahari, Pachwara Coalfield, Rajmahal Basin, Jharkhand, India. *Journal Palaeontological Society of India* 55: 29–35.

Tyson RV 1993. Palynofacies analysis; In Jenkins, D.G. (ed.): Applied Micropaleontology. Dordrecht, Kluwer 153-191.

Tyson RV 1995. Sedimentary Organic Matter: Organic Facies and Palynofacies. New York, Chapman & Hall, 615.

Utting J 1976. Pollen and spore assemblage in the Luwumbu Coal Formation (Lower Karoo) of the North Luangwa Valley, Zambia, and their biostratigraphic significance. Review of Palaeobotany and Palynology 21: 295–315.

Van Der Zwan CJ 1990. Palynostratigraphy and palynofacies reconstruction of the Upper Jurassic to lowermost Cretaceous of the Draugen Field, offshore mid-Norway. Review of Palaeobotany and Palynology 62: 157–186.

Vijaya 2006. Permian to Jurassic Palynoflora in Subsurface Strata, Pachami Area, West Bengal, India. Gondwana Geological Magazine 21: 13–26.

Vijaya 2009. Palynofloral changes in the Upper Palaeozoic and Mesozoic of the Deocha-Pachamari area, Birbhum Coalfield, West Bengal, India. Science in China Series D-Earth Sciences 52: 1932–1952.

Vijaya 2011. Palynostratigraphy of subsurface Upper Permian and Mesozoic Succession, Rakshitpur area, Raniganj Coalfield, West Bengal. Palaeoworld 20: 61–74.

Vijaya, Tripathi A, Roy A & Mitra S 2012. Palynostratigraphy and age correlation of subsurface strata within the Sub-basins in Singrauli Gondwana Basin, India. Journal of Earth System Science 121: 1071–1092.

Williams GL 1992. Palynology as a palaeoenvironmental indicator in the Brent Group, northern North Sea; *In*: Morton, A. C., Haszeldine, R. S., Giles, M. R. & Brown, S. (eds.), *Geology of the Brent Group*. Geological Society of London Special Publication 61: 203–212.

Wright RP & Askin RA 1987. The Permian-Triassic boundary in the southern Morondava Basin of Madagascar as defined by plant microfossils; *In*: McKenzie, G. D. (ed.), "Gondwana six: stratigraphy, sedimentology, and paleontology". Geophysical monograph 41: 157–166.

Zavattieri AM, Gutiérrez PR, Ezpeleta M & Astini RA 2008. Nuevos elementos de la microflora de la Formación La Veteada (Pérmico) en los afloramientos de la sierra de Narváez (Catamarca), Argentina. *Ameghiniana, Suplementos Resúmenes* 45: 10R–11R.

Zhang S, Tang S, Tang D, Pan Z & Yang F 2010. The characteristics of coal reservoir pores and coal facies in Liulin district, Hedong coal field of China. *International Journal of Coal Geology* 81: 117–127.

Appendix

Figure 1. Study area map of Godavari Graben with location map of the borehole in the Jangareddygudem area.

Figure 2. Lithological column and sample levels, stratigraphic occurrence, and relative abundance of various palynomorphs in borehole MJR-13.

Figure 3. Stratigraphically significant Palynotaxa of Late Permian

A. *Faunipollenites singrauliensis*, BSIP Slide No. 15328, P57/3

B. *Striatopodocarpites magnificus*, BSIP Slide No. 15329, K48/3

C. *Faunipollenites enigmatus*, BSIP Slide No. 15325, K49

D. *F. singrauliensis*, MJR-13, BSIP Slide No. 15328, U57/4

E. *Strotersporites* sp., BSIP Slide No. 15329, P45/4

F. *Strotersporites crassiletus*, BSIP Slide No. 15326, M44/3

G. *Striatites reticuloides*, BSIP Slide No. 15328, P32/4

H. *Striatopodocarpites* sp., BSIP Slide No. 15329, K55/2

I. *Striasulcites tectus*, BSIP Slide No. 15329, G62/1

J. *S. tectus*, BSIP Slide No. 15332, E62

K. *S. tectus*, BSIP Slide No. 15329, U54/4

L. *S. tectus*, BSIP Slide No. 15328, H38/4

M. *Faunipollenites singrauliensis*, BSIP Slide No. 15326, S33/3

N. *Faunipollenites* sp., BSIP Slide No. 15325, K35/2

O. *Crescentipollenites densus*, BSIP Slide No. 15327, H46

P. *Verticipollenites finimitus*, BSIP Slide No. 15322, H51/4

Figure 4. Stratigraphically significant Palynotaxa of Late Permian

A. *Densipollenites magnicarpus*, BSIP Slide No. 15321, W57/4

B. *D. invisus*, BSIP Slide No. 15325, J52

C. *D. invisus*, BSIP Slide No. 15325, J52/4

D. *D. invisus*, BSIP Slide No. 15325, K37/3

E. *D. indicus*, BSIP Slide No. 15320, E61/1

F. *D. marginalis*, BSIP Slide No. 15332, P53/4

G. *D. magnicarpus*, BSIP Slide No. 15331, O69/1

H. *Caheniasaccites elongatus*, BSIP Slide No. 15330, E63/1

I. *Scheuringipollenites maximus*, BSIP Slide No. 15329, Q46/2

J. *S. maximus*, BSIP Slide No. 15328, K67/4

K. *S. maximus*, BSIP Slide No. 15325, K51

L. *Cycadopites* sp., BSIP Slide No. 15329, F52/4

M. *Striasulcites* sp., BSIP Slide No. 15329, F52/4

N. *Falcisporites zapfei*, BSIP Slide No. 15332, W59/2

O. *Parasaccites densicarpus*, BSIP Slide No. 15324, M61/4

P. *P. diffusus*, BSIP Slide No. 15321, E50/2

Q. *Plicatipollenites ovatus*, BSIP Slide No. 15332, W55/3

R. *P. indicus*, BSIP Slide No. 15334, P64/4.

Figure 5. Stratigraphically significant Palynotaxa of Late Permian

A. *Striatopodocarpites multistriatus*, BSIP Slide No. 15325, G49/1

B. *S. multistriatus*, BSIP Slide No. 15325, M39/1

C. *Lunatisporites pellucidus*, BSIP Slide No. 15319, S61/2

D. *Striatites rhombicus*, BSIP Slide No. 15325, C37/4

E. *Guttulapollenites hannonicus*, BSIP Slide No. 15325, O63/4

F. *Corisaccites alutus*, BSIP Slide No. 15325, O36/4

G. *C. alutus*, BSIP Slide No. 15334, R60/2

H. *Praecolpatites nidpurensis*, BSIP Slide No. 15321, O68/4

I. *Lunatisporites* sp., BSIP Slide No. 15331, N69/4

J. *Horriditriletes ramosus*, BSIP Slide No. 15331, H49/3

K. *Latosporites colliensis*, BSIP Slide No. 15319, V52/4

L. *Maculatasporites* sp., BSIP Slide No. 15335, J61/1

M. *Strotersporites perfectus*, BSIP Slide No. 15326, H41/2

N. Unidentified bisaccate-monolete, BSIP Slide No. 15333, G53/1

O. *Faunipollenites singrauliensis*, BSIP Slide No. 15326, F43/3

P. *Vestigipollenited rudis*, BSIP Slide No.15326, Q36/2

Figure 6. Frequency distribution of organic matter and palynofacies identified in borehole MJR-13.

Figure 7. Representative photomicrographs of palynofacies components (all bars 50µm). A. Palynofacies showing palynomorphs and degraded terrestrial phytoclasts, BSIP Slide No. 15327, Q55/1, B. translucent wood, BSIP Slide No. 15329, K45/3, C. translucent wood, BSIP Slide No. 15328, L43/2, D. opaque wood phytoclasts, BSIP Slide No. 15329, J60/4, E. structured organic matter (leaf cuticle), BSIP Slide No. 15335, P63/2, F. Structured organic matter, BSIP Slide No. 15321, W49/1, G. degraded terrestrial matter, BSIP Slide No. 15333, M60/2, H. degraded terrestrial organic matter, BSIP Slide No. 15331, D45/2, I. Amorphous organic matter (AOM), BSIP Slide No. 15325, S41/1, J. Amorphous organic matter (AOM), BSIP Slide No. 15321, S45/1.

Figure 8. Ternary diagram showing the distribution of various palynofacies (after Aggarwal et al., 2019).

Figure 90. The number of palynotaxa recovered in borehole MJR-13 belongs to different plant groups.

Figure 10. Differential distribution of palynotaxa belonging to different plant groups at individual depth levels.

Table 1. *A generalized stratigraphical sequence in Chintalapudi Sub-basin (after Lakshminarayana, 1996).*

Table 2. *List of spores and pollen grains with their probable botanical affinities (Balme, 1995; Costamagna et al., 2018).*

ACCEPTED